- PI 572440 origin: United States. developed: D.E. Zimmer, A.L.
 Urie. origin institute: Agricultural Research Service -USDA, Crops Research Division, Logan, Utah 84321 United
 States. cultivar: PCM-1. pedigree: PI 195895/Frio, F3.
 other id: GP-2. source: Crop Sci. 10(4):463 1970.
 group: CSR-SAFFLOWER. other id: W6 12777. group: W6.
 remarks: Possess seedling and foliage rust-resistance
 superior to Ute. Equal in oil to Frio, 5 to 10cm shorter,
 and about a week earlier. In comparative yield trials in
 1969, yielded 5% more than US-10 (2,643kg/ha). Cultivar.
 Seed.
- PI 572441 origin: United States. developed: D.E. Zimmer, A.L.
 Urie. origin institute: Agricultural Research Service -USDA, Crops Research Division, Logan, Utah 84321 United
 States. cultivar: PCM-2. pedigree: PI 195895/U-5, F3.
 other id: GP-3. source: Crop Sci. 10(4):463 1970.
 group: CSR-SAFFLOWER. other id: W6 12778. group: W6.
 remarks: Possess seedling and foliage rust resistance
 superior to that of Ute. Lower in oil than Frio, 5 to
 10cm taller, and 5 days later. In comparative yield
 trials in 1969, outyielded Us-10 by 24% (3,114kg/ha).
 Cultivar. Seed.
- PI 572442 origin: United States. developed: D.E. Zimmer, A.L. Urie. origin institute: Agricultural Research Service --USDA, Crops Research Division, Logan, Utah 84321 United States. cultivar: PCN1. pedigree: Nebraska 1-1-5/Frio, other id: GP-4. source: Crop Sci. 10(4):463 1970. group: CSR-SAFFLOWER. other id: W6 7109. group: W6. remarks: Possesses the exclusionary rust resistance gene N1. This gene although effective against the seedling phase, is ineffective against the foliage phase. The utilization of a variety carrying the Nl gene would presumably greatly reduce primary inoculum for foliage infection, thereby reducing the need of foliage resistance. 5-7 days later in maturity than Frio, 5 to 10cm taller, and equal to Frio in oil percentage. Not been yield tested. Cultivar. Seed.
- PI 572443 origin: United States. developed: D.E. Zimmer, A.L.
 Urie. origin institute: Agricultural Research Service -USDA, Crops Research Division, Logan, Utah 84321 United
 States. cultivar: PCOY. pedigree: 59-270/Frio, F3.
 other id: GP-5. source: Crop Sci. 10(4):463 1970.
 group: CSR-SAFFLOWER. other id: W6 12779. group: W6.
 remarks: Highest type of foliage rust-resistance
 presently known in cultivated safflower; and its
 resistance to seedling rust is surpassed only by the
 exclusionary resistance of Nebraska 1-1-5. Produces seed
 equal in oil content to Frio and is of the same maturity
 and height. Has not been yield tested. Cultivar. Seed.